

<i>spironolactone/hctz</i>	25 / 25 mg	200/200 mg/day	Hyperkalemia, drowsiness, lethargy, mental confusion	1	1
<i>triamterene/hctz</i>	37.5 / 25 mg (1 or 2 tablets/ capsules) to 75mg/50mg (1 tablet daily)	75/50 mg/day	Hyperkalemia, dizziness, fatigue	1	1
Vasodilators					
<i>hydralazine</i>	75 mg	400 mg/day	Hypotension, palpitation, tachycardia, anorexia	2	1
<i>isosorbide</i>	30 mg	240 mg/day	Hypotension, headaches, dizziness, orthostatic hypotension, syncope	1	1

Brand-name drugs are capitalized and generic drugs are listed in lower-case italics

[NF] = Non-formulary [ST] = Step Therapy



CHF Guideline References

CHF Guideline adapted from:

- Circulation – Journal of the American Heart Association. Focused Update: ACCF/AHA Guidelines for the Diagnosis and Management of Heart Failure in Adults. A report of the American College of Cardiology Foundation / American Heart Association Task Force on Practice Guidelines. M. Jessup, WT. Abraham, DE. Casey, AM. Feldman, GS. Francis, TG. Ganiats, MA. Konstam, DM. Mancini, PS. Rahko, MA. Silver, LW. Stevenson and CW. Yancy. (2009, Mar 26). AHA: Dallas, TX.
- Drug Facts & Comparisons. (2010, Feb). Retrieved from www.online.factsandcomparisons.com. Wolters Kluwer Health: Deventer, The Netherlands